

Sigma-model solitons from the Schrödinger representation

Giovanni Landi

Trieste

Stochastic and Analytic Methods in Mathematical Physics

Yerevan, Armenia, September 4-11, 2016

Sigma-model solitons on noncommutative spaces

L. Dabrowski, [G.L. Franz Luef](#)
Lett. Math. Phys. 105 (2015) 1663–1688

and earlier work with L. Dabrowski and T. Krajewski
and more recent work by J. Rosenberg and Mathai V.

Abstract

Sigma-model solitons over the Moyal plane and noncommutative tori (also known as Irrational Rotation Algebra), as source spaces, with a target space made of two points

A natural action functional leads to self-duality equations for projections in the source algebra

Solutions, having non-trivial topological content (instantons), are constructed via suitable Morita duality bimodules (obtained using the Schrödinger representation).

Inputs from time-frequency analysis and Gabor analysis

Non-linear σ -models:

field theories of maps X between the **source** space (Σ, g) , and the **target** space (M, G) . The action functional

$$S[X] = \frac{1}{2\pi} \int_{\Sigma} \sqrt{g} g^{\mu\nu} G_{ij}(X) \partial_{\mu} X^i \partial_{\nu} X^j,$$

The stationary points:

harmonic maps from Σ to M ; describe minimal surfaces embedded in M .

Σ two dimensional, conformal invariance:

the action S is invariant for any rescaling of the metric $g \rightarrow e^{\sigma} g$.

Thus the action only depends on the conformal class of the metric and may be rewritten using a complex structure on Σ

$$S[X] = \frac{i}{\pi} \int_{\Sigma} G_{ij}(X) \partial X^i \wedge \bar{\partial} X^j,$$

Here ∂ and $\bar{\partial}$, a complex structure and $d = \partial + \bar{\partial}$.

In two dimensions

complex and conformal

are the same thing.

In two-dimensions, the conformal class of a general constant metric is parametrized by a complex number $\tau \in \mathbb{C}$, $\Im\tau > 0$.

Up to a conformal factor, the metric is

$$g = (g_{\mu\nu}) = \begin{pmatrix} 1 & \Re\tau \\ \Re\tau & |\tau|^2 \end{pmatrix}.$$

An algebraic generalization: by dualization and reformulation in terms of the *-algebras $\mathcal{A} = C^\infty(\Sigma, \mathbb{C})$ and $\mathcal{B} = C^\infty(M, \mathbb{C})$.

Embeddings X of Σ into M correspond to *-algebra morphisms $\pi_X : \mathcal{B} \rightarrow \mathcal{A}$, with correspondence

$$f \mapsto \pi_X(f) = f \circ X.$$

*-algebra morphisms make sense for general algebras \mathcal{A} and \mathcal{B} .

The configuration space is all *-algebra morphisms from \mathcal{B} to \mathcal{A}

The definition of the action functional involves generalizations of the conformal and Riemannian geometries.

Connes:

conformal is understood in the framework of positive Hochschild cohomology.

The tri-linear map $\phi : \mathcal{A}^{\otimes 3} \rightarrow \mathbb{R}$,

$$\phi(f_0, f_1, f_2) = \frac{i}{\pi} \int_{\Sigma} f_0 \partial f_1 \wedge \bar{\partial} f_2$$

is an extremal of positive Hochschild cocycles belonging to the Hochschild cohomology class of the cyclic cocycle ψ defined by

$$\psi(f_0, f_1, f_2) = \frac{i}{2\pi} \int_{\Sigma} f_0 df_1 \wedge df_2.$$

On the one hand ψ , the fundamental class, allows one to integrate 2-forms in dimension 2, so it is a **metric independent** object

On the other hand, ϕ defines a suitable positive scalar product

$$\langle a_0 da_1, b_0 db_1 \rangle = \phi(b_0^* a_0, a_1, b_1^*)$$

on 1-forms and **depends on the conformal class of the metric**.

Expressions like ϕ and ψ make sense for a general algebra \mathcal{A} .

Compose the cocycle ϕ with a morphism $\pi : \mathcal{B} \rightarrow \mathcal{A}$ to obtain a positive cocycle on \mathcal{B}

$$\phi_\pi = \phi \circ (\pi \otimes \pi \otimes \pi)$$

Evaluate the cocycle ϕ_π on a suitably element of $\mathcal{B}^{\otimes 3}$ which provides the noncommutative analogue of the metric on the target;

Easiest choice for this metric: a positive element $G = \sum_i b_0^i \delta b_1^i \delta b_2^i$ of the space of universal 2-forms $\Omega^2(\mathcal{B})$.

Thus, a well defined and positive quantity

$$S[\pi] = \phi_\pi(G) \tag{1}$$

a noncommutative analogue of the action functional of the non linear σ -model.

Here:

π is the dynamical variable (the embedding)

whereas ϕ (the conformal structure on the source) and G (the metric on the target) are background structures that have been fixed.

The critical points of the σ -model for the action functional (1): generalizations of harmonic maps; “minimally embedded surfaces” in the (noncommutative) space associated with \mathcal{B} .

The role of the other cocycle ψ is to give a topological ‘charge’.

More on this later on.

Two points as a target space $M = \{1, 2\}$

Any continuous map from a connected surface Σ to a discrete space is constant, a commutative theory would be trivial.

Not the case for a “noncommutative” source space: in general, not trivial such maps, ‘dually’, as $*$ -algebra morphisms from the algebra of functions over $M = \{1, 2\}$, that is \mathbb{C}^2 , to the algebra \mathcal{A} of the noncommutative source.

As a vector space \mathbb{C}^2 is generated by the projection function e defined by $e(1) = 1$ and $e(2) = 0$;

\Rightarrow any $*$ -algebra morphism $\pi : \mathbb{C}^2 \rightarrow \mathcal{A}$ is the same as a projection $p = \pi(e) \in \mathcal{A}$.

The configuration space of a two point target space sigma-model is the collection of all projections $P(\mathcal{A})$ in the algebra \mathcal{A} .

Choosing the metric $G = \delta e \delta e$ on the space $M = \{1, 2\}$, and a Hochschild cocycle ϕ for the conformal structure, the action functional is simply

$$S[p] = \phi(1, p, p),$$

Positivity in Hochschild cohom implies it is bounded by a **topological term**

Noncommutative torus and Moyal plane as source space \mathcal{A} :

the action functional is

$$S[p] = \frac{1}{4\pi} \text{tr}(\partial p \bar{\partial} p).$$

with the natural complex structure on \mathcal{A} given by

$$\partial = \partial_1 - i\partial_2, \quad \bar{\partial} = \partial_1 + i\partial_2,$$

and derivations ∂_1 and ∂_2 infinitesimal generators of a \mathbb{T}^2 -action

and tr an invariant trace.

All of above can be extended to more general metrics.

In two dimensions: Up to a conformal factor the general constant metric is parametrized by a complex number $\tau \in \mathbb{C}$, $\Im\tau > 0$.

The corresponding 'complex torus' $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z} + \tau\mathbb{Z}$ would act infinitesimally on \mathcal{A} with two complex derivations

$$\partial = \partial_1 + \bar{\tau}\partial_2, \quad \bar{\partial} = \partial_1 + \tau\partial_2.$$

As usual, the critical points of the action functional are obtained by equating to zero its first variation, that is the linear term in an infinitesimal variation

$$\delta S[p] = S[p + \delta p] - S[p], \quad \text{for } \delta p \in T_p(P(\mathcal{A})).$$

One gets

$$p \Delta(p) (1 - p) = 0 \quad \text{and} \quad (1 - p) \Delta(p) p = 0,$$

or, equivalently the non-linear equations of the second order

$$p \Delta(p) - \Delta(p) p = 0. \tag{2}$$

with the Laplacian of the metric $\Delta = \frac{1}{2}(\partial\bar{\partial} + \bar{\partial}\partial)$

The cyclic 2-cocycle giving the fundamental class is

$$\psi(a_0, a_1, a_2) = \frac{1}{2\pi i} \text{tr} (a_0(\partial_1 a_1 \partial_2 a_2 - \partial_2 a_1 \partial_1 a_2)),$$

For any projection $p \in P(\mathcal{A})$, the quantity

$$c_1(p) := \psi(p, p, p)$$

is an integer: the index of a Fredholm operator.

For any $p \in P(\mathcal{A})$ it holds that

$$S[p] \geq |c_1(p)|.$$

The equality for projection p satisfying **self-duality** or **anti-self duality** eqns

$$p(\partial_1 \pm i \partial_2)(p) = 0 \tag{3}$$

These equation imply the EOM (2).

Projection from **Morita equivalence** (Rieffel)

A Morita equivalence between (pre C^* -algebras) \mathcal{A} and \mathcal{B} :

a $\mathcal{A} - \mathcal{B}$ -bimodule \mathcal{E}

with a left-linear \mathcal{A} -valued **hp** $\bullet\langle \cdot, \cdot \rangle$ and a right-linear \mathcal{B} -valued **hp** $\langle \cdot, \cdot \rangle\bullet$.

There is an associativity condition:

$$\bullet\langle \xi, \eta \rangle \zeta = \xi \langle \eta, \zeta \rangle\bullet$$

It follows an identification $\mathcal{B} \simeq \mathcal{K}_{\mathcal{A}}(\mathcal{E})$ (compact endomorphisms).

In particular, there exist elements $\{\eta_1, \dots, \eta_n\}$ in \mathcal{E} such that

$$\sum_j \langle \eta_j, \eta_j \rangle\bullet = 1_{\mathcal{B}}.$$

Then, the associativity condition gives that the matrix $p = (p_{jk})$

$$p_{jk} = \bullet\langle \eta_j, \eta_k \rangle$$

is a projection in the matrix algebra $M_n(\mathcal{A})$.

Both algebras \mathcal{A} and \mathcal{B} are in the joint smooth domain of two commuting derivations ∂_1 and ∂_2 ; and have faithful invariant tracial states, which are compatible in the sense that:

$$\text{tr} \bullet \langle \xi, \eta \rangle = \text{tr} \langle \eta, \xi \rangle \bullet$$

Derivations are lifted to \mathcal{E} as (left and right) **covariant derivatives**:

$$\nabla_j : \mathcal{E} \rightarrow \mathcal{E}, \quad j = 1, 2,$$

$$\nabla_j(a \xi) = (\partial_j a) \xi + a (\nabla_j \xi) \quad \text{and} \quad \nabla_j(\xi b) = (\nabla_j \xi) b + \xi (\partial_j b)$$

compatible with both the \mathcal{A} -valued hermitian structure $\bullet \langle \cdot, \cdot \rangle$; the \mathcal{B} -valued hermitian structure $\langle \cdot, \cdot \rangle \bullet$

$$\partial_j(\bullet \langle \xi, \eta \rangle) = \bullet \langle \nabla_j \xi, \eta \rangle + \bullet \langle \xi, \nabla_j \eta \rangle$$

and

$$\partial_j(\langle \xi, \eta \rangle \bullet) = \langle \nabla_j \xi, \eta \rangle \bullet + \langle \xi, \nabla_j \eta \rangle \bullet$$

Lifting self-duality equations: **solitons**

The **holomorphic/anti-holomorphic**, connection on \mathcal{E} ,

$$\nabla = \nabla_1 - i\nabla_2, \quad \bar{\nabla} = \nabla_1 + i\nabla_2$$

lift to \mathcal{E} the complex derivations $\partial = \partial_1 - i\partial_2$ or $\bar{\partial} = \partial_1 + i\partial_2$.

The '**rank**' one case:

Seek solutions of the s-d eqs (3) of the form

$$p_\psi := \bullet\langle\psi, \psi\rangle \in \mathcal{A} \quad \text{with} \quad \psi \in \mathcal{E} \quad \text{such that} \quad \langle\psi, \psi\rangle_\bullet = \mathbf{1}_\mathcal{B}.$$

The projection p_ψ is a solution of the s-d eqs:

$$p_\psi \partial(p_\psi) = 0,$$

if and only if the vector ψ is a generalized eigenvector of $\bar{\nabla}$

i.e. there exists $\lambda \in \mathcal{B}$ such that

$$\bar{\nabla}\psi = \psi\lambda.$$

How to compute the topological charge:

The curvature of the covariant derivatives is defined as

$$F_{12} := \nabla_1 \nabla_2 - \nabla_2 \nabla_1$$

Let $\psi \in \mathcal{E}$ be such that $\langle \psi, \psi \rangle_{\bullet} = 1_B$ and $p_{\psi} := \bullet \langle \psi, \psi \rangle \in \mathcal{A}$ the corresponding projection. Then, its topological charge is:

$$c_1(p_{\psi}) = -\frac{1}{2\pi i} \operatorname{tr} \langle \psi, F_{12} \psi \rangle_{\bullet}.$$

Constant curvature: $F_{12} = -2\pi i q \operatorname{id}_{\mathcal{E}}$

the projection $p_{\psi} = \bullet \langle \psi, \psi \rangle$ has then topological charge

$$c_1(p) = q \operatorname{tr}(1_B) \in \mathbb{Z}$$

note that neither q nor $\operatorname{tr}(1_B)$ need be an integer

Moyal plane from **Schrödinger** representation

The projective representation of \mathbb{R}^2 on $L^2(\mathbb{R})$ defined for $\xi \in L^2(\mathbb{R})$ by

$$(\pi(z)\xi)(t) = e^{2\pi it\omega}\xi(t-x), \quad \text{for } z = (x, \omega). \quad (4)$$

$$\Rightarrow \quad \pi(z)\pi(z') = e^{-2\pi i x\omega'}\pi(z+z').$$

The map $c : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{T}$, $c(z, z') = e^{-2\pi i(x\omega')}$ is a 2-cocycle.

Its matrix-coefficients are defined for $\xi, \eta \in L^2(\mathbb{R})$ by

$$V_\eta\xi(z) := \langle \xi, \pi(z)\eta \rangle_{L^2(\mathbb{R})} = \int_{\mathbb{R}} \xi(t)\bar{\eta}(t-x)e^{-2\pi it\omega} dt$$

In signal analysis $V_\eta\xi$ is known as the short time Fourier transform

Moyal's identity: $\langle V_\eta \xi, V_\psi \varphi \rangle_{L^2(\mathbb{R}^2)} = \langle \xi, \varphi \rangle_{L^2(\mathbb{R})} \overline{\langle \eta, \psi \rangle_{L^2(\mathbb{R})}}$

An additional important consequence of this identity:

is a reconstruction formula for $\xi \in L^2(\mathbb{R})$ in terms of $\{\pi(z)\eta : z \in \mathbb{R}^2\}$.

Let η and ψ be in $L^2(\mathbb{R})$ such that $\langle \psi, \eta \rangle \neq 0$. Then for any $\xi \in L^2(\mathbb{R})$,

$$\xi = \langle \psi, \eta \rangle^{-1} \iint_{\mathbb{R}^2} \langle \xi, \pi(z)\eta \rangle \pi(z)\psi \, dz = \langle \psi, \eta \rangle^{-1} \iint_{\mathbb{R}^2} V_\eta \xi(z) \pi(z)\psi \, dz.$$

The twisted group algebra $L^1(\mathbb{R}^2, c)$ of \mathbb{R}^2 associated to the cocycle c .

For k and l in $L^1(\mathbb{R}^2)$, the twisted convolution $(k \natural l)$:

$$(k \natural l)(z) = \iint k(z')l(z - z')c(z', z - z') \, dz'$$

and twisted involution of $k \in L^1(\mathbb{R}^2)$:

$$k^*(z) = c(z, z) \overline{k(-z)} = e^{-2\pi i x \omega} \overline{k(-z)}$$

The integrated representation

$$K = \pi(k) = \iint_{\mathbb{R}^2} k(z)\pi(z)dz$$

for $k \in L^1(\mathbb{R}^2)$, is a non-degenerate bounded representation of the twisted convolution algebra $L^1(\mathbb{R}, c)$ on $L^2(\mathbb{R}^2)$.

The adjoint of $K = \pi(k)$ is given by $K^* = \pi(k^*)$ and the composition of $K = \pi(k)$ and $L = \pi(l)$ corresponds to $(k \natural l)$:

$$KL = \iint_{\mathbb{R}^2} (k \natural l)(z)\pi(z)dz.$$

Denote by \mathcal{A} the class of all operators $K = \pi(k)$ for $k \in \mathcal{S}(\mathbb{R}^2)$; they are all trace-class. Its norm closure is all compact operators.

\mathcal{A} is a model of the Moyal plane: the Fourier transforms of the symbols defining elements of \mathcal{A} yield the Moyal product:

$$k \star l = \mathcal{F}^{-1}(\mathcal{F}(k) \natural \mathcal{F}(l)) \quad \text{for } k, l \in \mathcal{S}(\mathbb{R}^2).$$

Rieffel:

(a result equivalent to the uniqueness of the Schrodinger representation of the CCR (for the two- dimensional quantum phase space of a free particle) :

The space $\mathcal{E} = \mathcal{S}(\mathbb{R})$ is an equivalence bimodule between \mathcal{A} and \mathbb{C}

with respect to the actions:

$$K \cdot \xi = \iint k(z) \pi(z) \xi \, dz,$$

$$\xi \cdot \lambda = \xi \bar{\lambda}$$

and \mathcal{A} and \mathbb{C} -valued hermitian products:

$$\bullet \langle \xi, \eta \rangle = \iint \langle \xi, \pi(z) \eta \rangle_{L^2(\mathbb{R})} \pi(z) \, dz = \iint V_\eta \xi(z) \pi(z) \, dz = \pi(V_\eta \xi)$$

$$\langle \xi, \eta \rangle_\bullet = \langle \eta, \xi \rangle_{L^2(\mathbb{R})}.$$

A two dimensional spectral geometry

Commuting derivations (an infinitesimal action of \mathbb{T}^2) ∂_1, ∂_2 :

$$\partial_1 K = 2\pi i \iint_{\mathbb{R}^2} x k(x, \omega) \pi(x, \omega) dx d\omega,$$

$$\partial_2 K = 2\pi i \iint_{\mathbb{R}^2} \omega k(x, \omega) \pi(x, \omega) dx d\omega.$$

They lift to covariant derivatives on the equivalence bimodule \mathcal{E} :

$$(\nabla_1 \xi)(t) = 2\pi i t \xi(t) \quad \text{and} \quad (\nabla_2 \xi)(t) = \xi'(t)$$

they are compatible with both left and right hermitian structures.

The connection has constant curvature:

$$F_{1,2} := [\nabla_1, \nabla_2] = -2\pi i \text{id}_{\mathcal{E}}$$

For $\psi \in \mathcal{S}(\mathbb{R})$ normalized as $\langle \psi, \psi \rangle_{\bullet} = \|\psi\|_2 = 1$,

\Rightarrow a non-trivial projection $p_{\psi} = \bullet \langle \psi, \psi \rangle$ in \mathcal{A} .

The projection p_{ψ} is a solution of the self-duality equations,

$$p_{\psi}(\partial p_{\psi}) = 0$$

if and only if, for some $\lambda \in \mathbb{C}$, the element ψ satisfies

$$\bar{\nabla} \psi = \psi \lambda.$$

Eigenfunction equations for $\bar{\nabla}$; solutions are generalized Gaussians:

$$\psi_{\lambda}(t) = c e^{-\pi t^2 - i\lambda t}.$$

Explicitly,

$$p_{\psi} = \bullet \langle \psi, \psi \rangle = \iint_{\mathbb{R}^2} V_{\psi} \psi(z) \pi(z) dz$$

$$V_{\psi} \psi(x, \omega) = e^{-\frac{\pi}{2}(x^2 + \omega^2)} e^{-\pi i x \omega - \frac{i}{2}(\bar{\lambda} + \lambda)x + \frac{i}{2}(\bar{\lambda} - \lambda)\omega}.$$

For its topological charge:

$$c_1(p_{\psi}) = \text{tr}(p_{\psi}) = V_{\psi} \psi(0) = 1.$$

The constant curvature is none other than the Heisenberg commutation relations (in the Schrödinger representation).

The anti-holomorphic connection $\bar{\nabla} = \nabla_1 + i\nabla_2$ is the annihilation operator; the holomorphic $\nabla = \nabla_1 - i\nabla_2$ is the creation operator.

The self-duality equation for the projections is the equation for the minimizers of the Heisenberg uncertainty relation, which explains why they are Gaussian ψ_λ .

The **irrational rotation algebra** (aka the noncommutative torus).

For $\theta \in \mathbb{R}$, the C^* -algebra A_θ of the noncommutative torus

is the norm closure of the span of $\{\pi(\theta k, l) : k, l \in \mathbb{Z}\}$: the restriction of the Schrödinger rep (4) of \mathbb{R}^2 on $L^2(\mathbb{R})$ to $\theta\mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$.

Denoting $\pi(0, 1) = M_1$ and $\pi(\theta, 0) = T_\theta$ they satisfy:

$$M_1 T_\theta = e^{2\pi i \theta} T_\theta M_1.$$

The smooth torus: subalgebra \mathcal{A}_θ of A_θ consisting of operators

$$\pi(\mathbf{a}) = \sum_{k, l \in \mathbb{Z}} a_{kl} \pi(\theta k, l), \quad \text{for } \mathbf{a} = (a_{kl}) \in \mathcal{S}(\mathbb{Z}^2).$$

With \mathbf{a} and \mathbf{b} in $\mathcal{S}(\mathbb{R})$ we have for their product

$$\pi(\mathbf{a})\pi(\mathbf{b}) = \pi(\mathbf{a} \sharp \mathbf{b})$$

where $\mathbf{a} \sharp \mathbf{b}$ is the twisted convolution

$$(\mathbf{a} \sharp \mathbf{b})(k, l) = \sum_{m, n \in \mathbb{Z}} a_{mn} b_{k-m, n-l} e^{-2\pi i \theta n(k-m)}$$

while $\pi(\mathbf{a})^* = \pi(\mathbf{a}^*)$, where \mathbf{a}^* is the twisted involution of \mathbf{a} :

$$(a_{kl})^* = e^{-2\pi i \theta kl} \overline{a_{-k, -l}}.$$

Operators commuting with $\pi(\theta k, l)$ are associated with the lattice $\mathbb{Z} \times \theta^{-1}\mathbb{Z}$.

They make up the algebra $\mathcal{A}_{1/\theta}$ of elements

$$b = \sum_{k, l \in \mathbb{Z}} b_{kl} \pi(k, \theta^{-1}l), \quad \text{for } \mathbf{b} = (b_{kl}) \in \mathcal{S}(\mathbb{Z}^2)$$

Take $\mathcal{A} = \mathcal{A}_\theta$ and $\mathcal{B} = (\mathcal{A}_{1/\theta})^{\text{op}} \simeq \mathcal{A}_{-1/\theta}$ (this acts from the right)

The space $\mathcal{E} = \mathcal{S}(\mathbb{R})$ is an equivalence bimodule between the noncommutative tori \mathcal{A} and \mathcal{B} with respect to the actions:

$$a \cdot \xi = \sum_{k,l \in \mathbb{Z}} a_{kl} \pi(\theta k, l) \xi,$$

and

$$\xi \cdot b = \sum_{k,l \in \mathbb{Z}} b_{kl} \pi(k, \theta^{-1} l)^* \xi,$$

and with hermitian products

$$\bullet \langle \xi, \eta \rangle = \theta \sum_{k,l \in \mathbb{Z}} V_\eta \xi(\theta k, l) \pi(\theta k, l),$$

and

$$\langle \xi, \eta \rangle \bullet = \sum_{k,l \in \mathbb{Z}} V_\xi \eta(k, l \theta^{-1}) \pi(k, \theta^{-1} l).$$

A two dimensional spectral geometry

The infinitesimal action of an ordinary torus \mathbb{T}^2 on both algebras \mathcal{A}_θ and $\mathcal{A}_{-1/\theta}$, are derivations. On \mathcal{A}_θ they are

$$\partial_1(a) = 2\pi i \sum_{k,l} k a_{k,l} \pi(\theta k, l)$$

and

$$\partial_2(a) = 2\pi i \sum_{k,l} l a_{k,l} \pi(\theta k, l),$$

and the dual ones on $\mathcal{A}_{-1/\theta}$ are then

$$\partial_1(b) = -2\pi i \theta^{-1} \sum_{k,l} k b_{k,l} \pi(k, \theta^{-1} l)^*$$

and

$$\partial_2(b) = -2\pi i \theta^{-1} \sum_{k,l} l b_{k,l} \pi(k, \theta^{-1} l)^*.$$

Lift to covariant derivatives ∇_1, ∇_2 on the bimodules $\mathcal{E} = \mathcal{S}(\mathbb{R})$:

$$(\nabla_1 \xi)(t) = 2\pi i \theta^{-1} t \xi(t) \quad \text{and} \quad (\nabla_2 \xi)(t) = \frac{d\xi(t)}{dt} =: \xi'(t).$$

The curvature is constant:

$$F_{1,2} := [\nabla_1, \nabla_2] = -2\pi i \theta^{-1} \text{id}_{\mathcal{E}}.$$

Frames

As a module over \mathcal{A}_θ , the space $\mathcal{E} = \mathcal{S}(\mathbb{R})$ is of finite rank and projective and it admits a **standard module Parseval frame** $\{\eta_1, \dots, \eta_n\}$ for $\mathcal{S}(\mathbb{R})$, that is each $\xi \in \mathcal{S}(\mathbb{R})$ has an expansion,

$$\xi = \bullet\langle \xi, \eta_1 \rangle \eta_1 + \cdots + \bullet\langle \xi, \eta_n \rangle \eta_n.$$

For $0 < \theta < 1$, the module $\mathcal{S}(\mathbb{R})$, is given by a projection in \mathcal{A}_θ itself: one can use a one-element Parseval frame η

From a standard module frame η one gets a Parseval frame $\tilde{\eta}$ by taking the element $\tilde{\eta} := \eta(\langle \eta, \eta \rangle_\bullet)^{-1/2}$

Then $\langle \tilde{\eta}, \tilde{\eta} \rangle_\bullet = 1$ and $\bullet\langle \tilde{\eta}, \tilde{\eta} \rangle$ is a projection in \mathcal{A}_θ .

Frames and projections:

- The Hermite function

$$\eta = \psi_k(t) = c_k e^{\pi t^2} \frac{d^k}{dt^k} e^{-2\pi t^2}$$

gives a projection $p_k = \bullet \langle \tilde{\eta}, \tilde{\eta} \rangle \in \mathcal{A}_\theta$, if $0 < \theta < (k + 1)^{-1}$.

- Let $\eta \in \mathcal{S}(\mathbb{R})$ be a totally positive function of finite type greater than 2. Then, $p_\eta = \bullet \langle \tilde{\eta}, \tilde{\eta} \rangle$ is a projection in \mathcal{A}_θ for $0 < \theta < 1$.

All of these projections have topological charge equal to 1. From

$$c_1(p) = q \operatorname{tr}(1_B)$$

with now $q = \theta^{-1}$ (the curvature) and $\operatorname{tr}(1_B) = \operatorname{tr}(\mathcal{A}_{-1/\theta}) = \theta$.

Duality and Gabor frames

For a Parseval frame, the *duality principle* (*Wexler-Raz identity*), reads as an expansion of each ξ in $\mathcal{S}(\mathbb{R})$ both over \mathcal{A} and \mathcal{B} ,

$$\xi = \bullet\langle\xi, \tilde{\eta}\rangle \tilde{\eta} = \tilde{\eta} \langle\tilde{\eta}, \xi\rangle\bullet,$$

with $\bullet\langle\xi, \tilde{\eta}\rangle \in \mathcal{A}$ and $\langle\tilde{\eta}, \xi\rangle\bullet \in \mathcal{B}$ which are uniquely determined.

This helps for the soliton equation.

As before, the s-d eqs for the projection p_ψ obeys $p_\psi \partial(p_\psi) = 0$ translate to a generalized eigenvector equation

$$\bar{\nabla}\psi = \psi\lambda,$$

with now $\lambda = \langle \psi, \bar{\nabla}\psi \rangle_\bullet \in \mathcal{A}_{-1/\theta}$.

Using the duality principle we have that

$$\text{with } \psi := \eta(\langle \eta, \eta \rangle_\bullet)^{-1/2}$$

the projection $p_\psi = \bullet\langle \psi, \psi \rangle \in \mathcal{A}_\theta$

satisfies the s-d eqs:

- For $0 < \theta < (k + 1)^{-1}$, if η is the k -th Hermite functions ψ_k .
- For $0 < \theta < 1$, if η is a tot pos fun in $\mathcal{S}(\mathbb{R})$ of finite type greater than 2.

In particular, the Gaussian function

$$\psi_\lambda(t) = c e^{-\frac{\pi}{\theta} t^2 - i\lambda t}, \quad \text{for } \lambda \in \mathbb{C},$$

obeys the equation $\bar{\nabla}\psi_\lambda = \psi_\lambda\lambda$.

The right hermitian product $\langle \psi_\lambda, \psi_\lambda \rangle_\bullet$ is invertible in $\mathcal{A}_{-1/\theta}$ for all $0 < \theta < 1$,

so that the projections $p_\lambda = \bullet \langle \tilde{\psi}_\lambda, \tilde{\psi}_\lambda \rangle$, with $\tilde{\psi}_\lambda := \psi_\lambda (\langle \psi_\lambda, \psi_\lambda \rangle_\bullet)^{-1/2}$

are solutions of the self-duality equations

The moduli space of such Gaussian solutions, is parametrised by possible λ 's modulo gauge transformations

(implemented by invertible elements in $\mathcal{A}_{-1/\theta}$)

is a copy of the complex torus.

More examples and applications

Sigma-model solitons over the Moyal plane and noncommutative tori,
as source spaces, with a target space made of two points

A natural action functional leads to self-duality equations for projections in
the source algebra

Solutions, having non-trivial topological content, are constructed via suitable
Morita duality bimodules,

Inputs from time-frequency analysis and Gabor analysis

More interesting cases

Uses in time-frequency analysis and Gabor analysis coming up

Thank you